

Biomechanical investigation of dynamic topping-off in two-level posterior instrumentation

*W. Schmoelz¹, A. Keiler¹, S. Hartmann², K.-M. Scheufler², C. Thomé², A.A. Hegewald²

¹Medizinische Universität Innsbruck, Unfallchirurgie, Innsbruck, Österreich

²Medizinische Universität Innsbruck, Neurochirurgie, Innsbruck, Österreich

Introduction:

Topping-off instrumented lumbar fusion with a dynamic implant has been suggested to avoid adjacent segment disease by creating a smoother transition zone from the instrumented segments to the untreated levels above. This study evaluates ex-vivo the range of motion (RoM) and intradiscal pressure to characterize the transition zones of two-level posterior instrumentation strategies for elucidating biomechanical differences between rigid fixation and the topping off approach with a pedicle screw-based dynamic implant.

Material and Methods:

Eight human lumbar spines (L1-5) were loaded in a spine tester with pure moments of 7.5Nm in lateral bending (LB), flexion/extension (FE) and axial rotation (AR) and with a hybrid loading protocol in lateral bending and flexion/extension. The following states were tested: (a) intact, (b) laminectomy L4 with rigid fixation of L4-5 and dynamic topping off L3-4 (HPS, Paradigm Spine, Wurmlingen, Germany). (c) laminectomy L4 with rigid fixation of L3-5. The RoM for all segments for both loading protocols was evaluated and normalized to the intact segmental RoM. The intradiscal pressure was measured in the segments L3-4 and L2-3.

Results:

For pure moment loading, RoMs of the segments cranial to both instrumentations (L1-2 and L2-3) were not affected by the type of instrumentation ($p > 0.5$) in FE, LB and AR. The dynamic instrumentation in L3-4 reduced the RoM compared to intact ($p < 0.05$) but allowed more motion than the rigid fixation of the same segment ($p < 0.05$) in LB and FE. Under the hybrid loading protocol, the cranial segments (L1-2 and L2-3) had a significant higher RoM ($p < 0.05$) for both instrumentations compared to the intact (ranging from 162%-227% of the intact RoM in FE and LB). Comparing the two instrumentation approaches with each other, the dynamic topping off showed a smaller increase of RoM than the rigid fixation (L1-2 $\Delta = 15\%$ and 26% ; L2-3 $\Delta = 8\%$ and 11% for FE and LB, respectively). Under pure moment loading the intradiscal pressure for FE and LB showed comparable effects for the untreated segment L2-3 (fig. 1) and comparable pressure reductions for both instrumentations in L3-4 (fig. 2).

Conclusion:

Regardless of the approach, two-level posterior instrumentation was accompanied by a considerable amount of compensatory movement in the cranial untreated segments under the hybrid protocol. Dynamic topping-off, however, showed a significant reduction of this compensatory movement in comparison to rigid fixation along with more than 50% of movement preservation of the dynamically instrumented level.

The clinical relevance of these differences cannot be deduced from this model. From a biomechanical point of view, however, they could offer advantages for the dynamic topping-off because of a slightly smoother transition zone.

Figure 1:




Figure 2:

